

schülke -†



Antiseptic for the treatment of vaginal infections

octenisept® vaginal therapeutic agent

Our plus:

- treats the symptoms of itching, burning and discharge due to bacteria in the vaginal area
- optimal moistening of the mucous membrane
- no restricted application for pregnant women after the 3rd month
- colourless

Application areas

- For the treatment of vaginal infections
- For the alleviation of symptoms in case of itching, burning and discharge due to bacteria in the vaginal area

Application methods

Insert the vaginal applicator deeply into the vagina, pressing down the actuator cap of the applicator ten times one by one, in the morning and evening on the first day of treatment, then once a day in the evening. The period of treatment is seven days. Prior to first application, remove the protection and actuator cap from the bottle and keep it in a safe place, then take the applicator from the wrapping and place it on the bottle.

It is recommended to insert the applicator deep into the vagina while sitting on the toilet or lying in bed. Then, the cap of the applicator must be pressed down ten times one by one. After application, the applicator should be wiped off externally and some preparation should be sprayed out with the applicator facing down.

Before next application, some preparation should be sprayed out once or twice with the applicator facing down.

In order to prevent a renewed infection, the partner should be involved in the treatment. For this purpose, the preparation is to be sprayed, without applicator and with actuator cap placed on top, on the male genital until it is completely moistened.

If the symptoms persist and in case of relapse, a gynaecological examination incl. microbiologic diagnostics and, if required, antibiotic or antifungal therapy, is necessary.

Microbiological efficacy:

octenisept® vaginal therapeutic agent is effective against:

- bacteria including chlamydia and mycoplasmas
- fungi and yeasts
- protozoa (trichomonads)
- virus (herpes simplex, HBV, HCV and HIV)

Product data

Composition:

100 g solution contains: pharmaceutically effective ingredients: 0.1g octenidindihydrochloride, 2.0g phenoxyethanol (Ph.Eur.)

Other ingredients: (3-coco fatty acid amidopropyl)-dimethylzaniumylacetate, sodium-D-gluconate, glycerol 85%, sodium chloride, purified water, sodium hydroxide

Chemisch-physikalische Daten:

Appearance:	colourless
Flash point:	> 99° C
pH value:	6.0 ± 0.5

octenisept® vaginal therapeutic agent

Special indications

- octenisept® vaginal therapeutic agent should not be used to rinse the abdominal cavity (e.g. during operation), the urinary bladder and the nose and nor be applied to the eardrum.
- Do not swallow octenisept® vaginal therapeutic agent in larger quantities, nor let it enter the bloodstream in larger quantities, e.g. by means of an inadvertent injection.
- Do not mix octenisept® vaginal therapeutic agent with other preparations.
- When applying octenisept® vaginal therapeutic agent, a momentary burning can occur as a subjective symptom in rare cases.
- octenisept® vaginal therapeutic agent shall not be used after the expiration date.
After the first use: Dispose the residual solution remaining in the container after termination of the treatment.
- octenisept® vaginal therapeutic agent is to be kept out of reach of children

Delivery forms / packaging units

These products are not available in every country. For any information please contact our local subsidiary or distributor.

Container sizes	Pack units	Art. No.
50 ml bottle with applicator	20 bottles per carton	121441

Environmental information

schülke manufactures products economically and with advanced, safe and environmentally friendly production processes, while at the same time maintaining our high quality standards.

Expert Opinion and Information

For an overview of our products containing Octenidine, please refer to our website www.schuelke.com

Should you have any further questions:

Customer Care

Phone: +49 (0)40 521 00-666

E-mail: info@schuelke.com

octenisept® vaginal therapeutic agent

- **Composition:** active pharmaceutical ingredients: 100g of solution contain: octenidine dihydrochloride 0.1g, phenoxyethanol (Ph.Eur.) 2.0g. Other ingredients: (3-coconut fatty acid amidopropyl)-dimethylazaniumylacetate, sodium-D-gluconate, glycerol 85%, sodium chloride, sodium hydroxide, purified water.
- **Application areas:** For the alleviation of symptoms in case of itching, burning and discharge due to bacteria in the vaginal area.
- **Contraindications:** In case of hypersensitivity to one or more of the active ingredients of the pharmaceutical. No sufficient experience is available regarding the application in children under the age of 8. octenisept® vaginal therapeutic agent should not be used during the first three months of pregnancy as a special precaution.
- **Adverse effects:** In rare cases, burning and itching in the vagina can occur after the application of octenisept®. Should you notice adverse effects which are not mentioned in this user information, please inform your doctor or pharmacist.
- **Warning notices and special preventive measures during application:** Do not swallow octenisept® vaginal therapeutic agent in larger quantities, nor let it enter the bloodstream in larger quantities.
- **Pharmaceutical company:**
Schülke & Mayr GmbH, 22840 Norderstedt, Germany

- V01 -



Schülke & Mayr GmbH is certified according to DIN EN ISO 9001, DIN EN ISO 14001 and DIN EN ISO 13485 (Reg.-No. 004567-MP23) and has a validated environmental system in accordance with the Eco Audit Regulation (Reg.-No. DE-150-00003).



Schülke & Mayr GmbH
22840 Norderstedt, Germany
Phone +49 (0) 40 - 521 00 - 0
Fax +49 (0) 40 - 521 00 - 318
www.schuelke.com
mail@schuelke.com

Schülke & Mayr UK Ltd.
1 Jenkin Road
GB-Sheffield S9 1AT
Phone +44 (0) 1142 - 54 35 - 00
Fax +44 (0) 1142 - 54 35 - 01
mail.uk@schuelke.com

Schülke & Mayr Ges.m.b.H
Seidengasse 9
1070 Wien, Austria
Phone +43 (0) 1 - 523 25 01 - 0
Fax +43 (0) 1 - 523 25 01 - 60
office@schuelke.at