

# schülke -†



Aqueous wound and mucous membrane antiseptic

## octenisept®

### Our plus:

- broad spectrum of antiseptic efficacy
- very good skin and mucous membrane compatibility
- suitable for infants and premature babies
- also safe use during pregnancy\*
- painless application
- colourless

### Application areas

- Wound treatment: Antiseptic treatment of traumatic, acute, chronic, surgical and burn wounds.
- Mucous membrane antiseptic:
  - Prior to diagnostic and surgical interventions in the anogenital and the urogenital area (e. g. before placing an intra-uterine device (IUD), before prenatal, intranatal and postnatal manipulations or before obliteration of haemorrhoids)
  - Prior to diagnostic and surgical interventions in the oral area (e.g. before tooth extractions or curettages)
  - Before placing transurethral disposable and indwelling catheters
  - For preoperative skin antisepsis in the area close to mucous membranes (e. g. before section)
  - MRSA: For antimicrobial whole body wash with MRSA colonised patients.

### Microbiological efficacy

octenisept® is effective against · bacteria including chlamydia and mycoplasma · fungi and yeasts · protozoa (trichomonads) · viruses (Herpes simplex, HBV, HCV and HIV)

### Contact times

Microbes	Contact time*
Bacteria - incl. the wound bacteria MRSA, E. coli, Pseudomonas aeruginosa - incl. chlamydia, Mycoplasma hominis, Ureaplasma urealyticum, Gardnerella vaginalis and Neisseria gonorrhoea	30 sec.
Fungi (Candida albicans)	2 min.
Malassezia furfur	1 min.
Protozoa (trichomonads)	1 min.
Viruses herpes simplex, HBV, HCV, HIV	30 sec.

Application area	Contact time*
Anogenital area	1 min.
Before catheterising the urinary bladder	1 min.
Skin antiseptic pre-section	at least 2 min.
Oral cavity Swab method Rinsing	2 min. 1 min.
Wounds	at least 1 min.
Whole body wash with MRSA patients	50% solution / 2 min.

\*Details on dosing, duration and way of application are available in the instructions for use.

\* Briese et al. (2010): Efficacy and tolerability of a local acting antiseptic agent in the treatment of vaginal dysbiosis during pregnancy; in Arch Gynecol Obstet

# octenisept®

## Application methods / indications

- The area of the mucous membrane and skin to be treated must be moistened evenly and thoroughly with the antiseptic.
- Swab method: Under aseptic conditions the areas of mucous membrane and skin should be rubbed with saturated swabs. Due to its anatomical and physiological state this method is preferred on the mucous membrane.
- Spray method: In individual cases, octenisept® may also be sprayed directly on accessible areas of the skin and mucous membrane. Make sure all areas are evenly moistened.
- Rinsing: octenisept® can also be used for oral and vaginal rinsing.

## Product data

### Composition:

100 g solution contains: Octenidine dihydrochloride 0.1 g, 2-Phenoxyethanol (Ph.Eur.) 2.0 g

### Chemical-physical data:

Appearance:	colourless
Flashpoint:	> 99°C
pH value:	6.0 ± 0,5

## Special advice

- When irrigating wounds, ensure that the preparation is not inserted under pressure or injected into the tissue.
- When rinsing wound cavities make sure that there is always a drainage point (e.g. drain, tab).
- octenisept® should not be used for rinsing the abdominal cavity (e.g. intraoperatively), the urinary bladder and nose or eardrum.
- Do not swallow octenisept® or allow it to reach the bloodstream, e.g. by being accidentally injected.
- Do not mix octenisept® with other compounds.
- Do not use octenisept® with PVP-iodine based antiseptics on adjacent areas of the skin, as it may lead to strong brown to violet discolouration.
- Bandages and incision foils can be applied once octenisept® has dried completely.
- In rare cases octenisept® may cause a temporary slight burning sensation.
- octenisept® can be warmed to body temperature.
- octenisept® should not be used after the expiry date.
- Once the container has been opened octenisept® should be used for no longer than three years but not beyond the expiry date.

## Delivery form / packaging units

These products are not available in every country. For more information please contact our local subsidiary or distributor.

Container sizes	Pack units	Art. No.
<b>Octenisept® Antiseptic Pre-catheterising of the bladder</b> 15 ml bottle	100 x 15 ml	121407
<b>Octenisept® Wound disinfection</b> 50 ml bottle with overhead spray pump	20 x 50 ml	121418
<b>Octenisept®</b> 250 ml bottle	10 x 250 ml	121406
250 ml bottle with overhead spray pump	10 x 250 ml	121411
500 ml bottle	20 x 500 ml	121464
1 l bottle	10 x 1 l	121403

## Environmental information

schülke manufactures products economically and with advanced, safe and environmentally friendly production processes, while at the same time maintaining our high quality standards.

## Expert opinions and information

For an overview of our products containing Octenidine, please refer to our website [www.schuelke.com](http://www.schuelke.com)

Should you have any further questions:

Customer Care

Phone: +49 (0)40 521 00-666

E-mail: [info@schuelke.com](mailto:info@schuelke.com)

**octenisept® • Composition:** 100 g solution contain: octenidine hydrochloride 0.1 g, phenoxyethanol (Ph.Eur.) 2.0 g. Other ingredients: (3-amidopropyl cocoate) dimethylammonium acetate, sodium D gluconate, glycerol 85%, sodium chloride, sodium hydroxide, purified water. • **Indications:** For repeated, short-term antiseptic treatment of mucous membranes, adjacent skin and as adjuvant antiseptic wound treatment. • **Contraindications:** octenisept® may not be used in cases of hypersensitivity to any of the components of the preparation. octenisept® should not be used for rinsing the abdominal cavity (e.g. intra-operatively) or the bladder, nor the tympanic membrane. • **Special warnings and special precautions for use:** Do not swallow octenisept® and do not allow octenisept® to pass into the circulation, e.g. as a result of accidental injection. When irrigating wounds, ensure that the preparation is not inserted under pressure or injected into the tissue. When irrigating wound cavities, ensure that there is always a drainage point (e.g. drain, tab). • **Undesirable effects:** In rare cases transient signs of local intolerance such as a slight burning sensation, redness or itching may occur at the application site. Rinsing of the oral cavity may cause a transitory bitter sensation. • **Manufacturer:** Schülke & Mayr GmbH, 22840 Norderstedt, Germany

If you notice any side effects not mentioned in this information, please inform your doctor or pharmacist.



Schülke & Mayr GmbH is certified according to DIN EN ISO 9001, DIN EN ISO 14001 and DIN EN ISO 13485 (Reg.-No. 004567-MP23) and has a validated environmental system in accordance with the Eco Audit Regulation (Reg.-No. DE-150-00003).



**Schülke & Mayr GmbH**  
22840 Norderstedt, Germany  
Phone +49 (0) 40 - 521 00 - 0  
Fax +49 (0) 40 - 521 00 - 318  
[www.schuelke.com](http://www.schuelke.com)  
[mail@schuelke.com](mailto:mail@schuelke.com)

**Schülke & Mayr UK Ltd.**  
1 Jenkin Road  
GB-Sheffield S9 1AT  
Phone +44 (0) 1142 - 54 35 - 00  
Fax +44 (0) 1142 - 54 35 - 01  
[mail.uk@schuelke.com](mailto:mail.uk@schuelke.com)

**Schülke & Mayr Ges.m.b.H**  
Seidengasse 9  
1070 Wien, Austria  
Phone +43 (0) 1 - 523 25 01 - 0  
Fax +43 (0) 1 - 523 25 01 - 0  
[office.austria@schuelke.com](mailto:office.austria@schuelke.com)