schülke

acts in 30 seconds

Antiseptic mouthrinse effectively combats the root causes of gingivitis and periodontitis.

octenident® antiseptic

1 mg/ml oromucosal solution Octenidine dihydrochloride

Our Plus

- Broad spectrum of activity inc. gram-negative bacteria
- Acts in 30 seconds
- Compliance with VAP guidelines in accordance with KRINKO
- Low irritation/allergic potential
- Alcohol-free
- Peppermint flavoured
- Can also be used during pregnancy*

*Please read the package leaflet.

Application area

octenident® antiseptic contains the active substance octenidine dihydrochloride.
octenident® antiseptic has an antimicrobial effect. It is used for

- temporary reduction of bacterial count in the oral cavity
- for temporary inhibition of plaque formation
- in cases of insufficient oral hygiene capacity (no toothbrushing possible, for example) in adults.

Instructions for use

- Use the solution twice a day (after meals and toothbrushing)
- Fill 10 ml in the measuring cup provided
- If you use octenident® antiseptic just after brushing, rinse your mouth thoroughly with water before rinsing with octenident® antiseptic
- Rinse your mouth thoroughly for about 30 seconds and spit out.
- The solution should not be swallowed
- Do not rinse the mouth with water directly afterwards
- Do not use this medicine for more than five days in a row

Microbiological efficacy

	Test method	Contact time
Bacteria, Yeast and Mycobacteria		
bactericidal	EN 13727	30 sec.
yeasticidal	EN 13624	1 min.

Product tested under clean conditions.

Product data

Composition:
100 g solution contains:
0.1 g Octenidine dihydrochloride

Other components: Glycerol, Sodium D-Gluconate, Citric Acid, Disodium Phosphate, PEG-40 Hydrogenated Castor Oil, Sucralose, Purified Water, Aroma

Chemical-physical data

Color	almost colourless
Density	ca. 1,0 g/cm³ / 20 °C
Flash point	> 100 °C / Method: ISO 2719
Form	liquid
pH	ca. 6,0 / 20 °C
Viscosity, dynamic	ca. 1 mPa*s

octenident® antiseptic

1 mg/ml oromucosal solution Octenidine dihydrochloride

Special advice

octenident® antiseptic is intended for superficial use only. It must not be administered deep in the tissue for example with a syringe or a blunt cannula in order to prevent irritations of the surrounding tissues potentially ranging from slight swellings to serious tissue necrosis. Specifically, it should not be used for the rinsing of periodontal pockets, of root canals or of wound cavities following dental surgery.

Information for order

Item	Delivery form	Item no.
octenident® antiseptic 250 ml bottle	12 / carton(s)	on request

These products are not available in every country. For more information please contact our local subsidiary or distributor.

Related products

- **octenident®** mouthrinse

Environmental information

schülke manufactures its products using advanced and safe production processes while maintaining high quality standards.

Expert opinion and information

Please visit our website for an overview of all available literature/reports on the product:

www.schuelke.com.

For individual questions:

Application Department

Phone: +49 40 52100-666

E-Mail: info@schuelke.com

octenident® antiseptic 1 mg/ml oromucosal solution
Active substance: octenidine dihydrochloride. **Composition:** 1 ml of solution contains 1 mg of octenidine dihydrochloride. Other ingredients: glycerol 85 per cent (E 422), sodium gluconate, citric acid, disodium phosphate dihydrate (for pH adjustment), macrogolglycerol hydroxystearate, sucralose, purifed water, mint flavour (contains propylene glycol (E 1520)).
Indications: octenident antiseptic has an antimicrobial effect. It is used for temporary reduction of bacterial count in the oral cavity, for temporary inhibition of plaque formation, and in cases of insufficient oral hygiene capacity (no toothbrushing possible, for example) in adults.
Contraindications: Allergy to octenidine dihydrochloride or any of the other ingredients.
Undesirable effects: Very common: Temporary taste disturbance, such as bitter aftertaste; Mild, reversible dental discoloration. Common: Numb sensation in the mouth, Coating of the mouth or the tongue, Temporary tongue discoloration, Sensitivity of teeth. Uncommon: Headache, Nausea, Tingling of the tongue, More saliva in the mouth than normal. Revision 09/21 If any of the side effects get serious, or if you notice any side effects not listed in this user information, please tell your doctor or pharmacist. Schülke & Mayr GmbH, 22840 Norderstedt, Germany, Tel. +49 40 52100-666, info@schuelke.com

schülke

schülke Headquarters
Schülke & Mayr GmbH
Robert-Koch-Straße 2
22851 Norderstedt | Germany
Phone +49 40 52100-666
www.schuelke.com
info@schuelke.com

Schülke & Mayr GmbH holds a Manufacturer's Authorisation according to sect 13 para 1 German Drug Law and Certificates of GMP Compliance for medicinal products.

/ 913531-ZZ-en-ZZ-02.12 06.2026
This product information is not automatically updated.